

The Chemical Evolution of Milky Way Satellite Galaxies

from Keck/DEIMOS Multi-Element Abundance Measurements

Evan Kirby (Caltech)

Abstract. A Keck/DEIMOS spectroscopic campaign of eight Milky Way (MW) dwarf spheroidal (dSph) satellite galaxies has generated spectral synthesis-based abundance measurements for nearly 3000 stars. The elements measured are Fe and the α elements Mg, Si, Ca, and Ti. The dSph metallicity distributions show that the histories of the less luminous dSphs were marked by massive amounts of gas loss. The $[\alpha/\text{Fe}]$ distributions indicate that the early star formation histories of most dSphs were very similar and that Type Ia supernova ejecta contributed to the abundances of all but the most metal-poor ($[\text{Fe}/\text{H}] < -2.5$) stars. Finally, a numerical chemical evolution model reveals that the star formation history of a dSph is a strong function of its present-day luminosity, but not velocity, dispersion, half-light radius, or Galactocentric distance.




Figure 1. Metallicity distributions of MW dSphs, ordered from most to least luminous. As dSph luminosity decreases, the average $[\text{Fe}/\text{H}]$ also decreases, a sign that the less luminous dSphs lost a great deal of gas during their star formation histories. Furthermore, the more luminous dSphs fit an analytic chemical evolution model with gas inflow better than the less luminous dSphs. The metallicity distributions of the less luminous dSphs are less peaked, more symmetric, and better matched to a leaky box model.




Figure 2. Moving averages of the $[\alpha/\text{Fe}]$ ratios in MW dSphs. The figure legend lists the dSphs in order of decreasing luminosity. At $[\text{Fe}/\text{H}] < -1.2$, the $[\alpha/\text{Fe}]$ ratios follow nearly the same path in all dSphs, suggesting similar star formation histories at early times. Furthermore, with few exceptions, there are no $[\alpha/\text{Fe}]$ plateaus at $[\text{Fe}/\text{H}] > -2.5$, which indicates that Type Ia supernova ejecta contributed to all but the most metal-poor stars.




Figure 3. Top left: Observed abundance distributions for the Sculptor dSph (black) with a best-fit chemical evolution model (red) that tracks the star formation rate, Type II and Ia supernova element production, and supernova feedback. Top right: Gas inflow rate, mass, star formation rate, metallicity, and $[\text{Mg}/\text{Fe}]$ as a function of time in the model. Bottom: Observed and modeled abundance distributions for the more luminous dSph Leo I and the fainter dSph Draco.

Additional reading:
 Kirby et al. 2010, ApJS, 191, 352, arXiv:1011.4516
 Kirby et al. 2011, ApJ, 727, 78, arXiv:1011.4637
 Kirby et al. 2011, ApJ, 727, 79, arXiv:1011.5221

Acknowledgments: This work is based in part on data obtained by the Keck Observatory, which is operated by the University of California, Lick Observatory, and the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26565.

Background image of the Fornax dSph: European Southern Observatory/Digital Sky Survey 2.